

My Account

Login
Create Account

Resources

View All (813)
Adenoviruses (137)
Antibodies (175)
Bioimages (67)
Genomics Studies (145)
mESC Lines (68)
Mouse Strains (120)
Miscellaneous (46)
Protocols (55)
Research Data (4)
Resource Tags (389)
Visualization (9)

Research & Cores

Core Facilities (5)
Research Highlights (5)
Research Networks
Research Objectives

Information


About the BCBC
BCBC Events
Branding & Logos
Career Opportunities
Health
NIH hESC Registry
Policies & Guidelines
Member Publications
Research Programs
Research Investigators
Member Directory
Tutorials

Nkx2.2^{LCA} - ES Cell Line RES257**ESC Line Information**

Cell Line Name:	Nkx2.2 ^{LCA}
Parental Cell Line:	TL-1
Background Strain:	129
Culturing Protocol:	Std_mESC_Culture.doc
Description:	This ES cell line contains a loxed cassette acceptor (LCA) allele that may be utilized for the exchange of DNAs of interest into the Nkx2.2 locus by recombinase mediated cassette exchange (RMCE). This LCA uses a Lox71/66 and Lox2272 strategy for RMCE and allows for manipulation of a 5.1 kb region of the gene (contains promoter, exon 1, and exon 2). This leads to a variety of experiments that can be performed using a wide range of cassette designs.

Genetic Alterations**1) Targeted Mutagenesis**

Type of Allele	Cassette Acceptor
Targeted Gene	NK2 transcription factor related, locus 2 (Nkx2-2 - NCBI GeneID:18088)
Targeted Allele	targeted mutation 1 (Nkx2.2 ^{tm1(LCA)})
Description of Targeting Vector	Through homologous recombination in ES cells, a 5.115 kb region of the Nkx2.2 gene is replaced by a floxed tk-neo cassette, a puromycin-thymidine kinase fusion gene driven by the mouse phosphoglycerol kinase promoter (pUTK) and a neomycin resistant gene driven by the bacterial EM7 promoter (EM7neo) flanked by minimal (34 bp) tandemly oriented lox71 and lox2272 sites (Cre-recombinase recognition sequences).
Targeting Vector Genbank File	pNkx2.2.TV.gb
Citations	Not Available

Associated Images**Image 1****Description:**

Through homologous recombination in ES cells, a 5.115 kb region of the Nkx2.2 gene was replaced with a pgk-puromycin-delta thymidine kinase [pu(Delta)TK]-fusion gene and a EM7-Kanamycin resistance cassette flanked by tandemly oriented lox71 and lox2272 sites (Cre-recombinase recognition sequences).


Reference:

Not provided


Repositories

Magnuson Lab

Access Status

 This resource is publicly viewable.

Request this Resource

 Request from a repository


Primary contributor: [Magnuson Lab](#)


Co-contributed by:

- [BCBC Mouse / ES Cell Core](#)

Resource Tags

embryonic, es, ESC, esc, LCA, mESC Core, Nkx2.2, Nkx2.2^{LCA}, RMCE, stem, TL-1

 Login to edit tags

 Read more about tags

Resource History & Actions

Approved on Mar 25, 2008

Last modified on Mar 23, 2015

 Login to edit or request an edit


Related resources**BCBC**

No matching resources

Other Consortia

No matching resources

Data courtesy of [dkCOIN](#). Only public resources are displayed.

 Request this resource

Stock #: *Not provided*

Availability Notes: *Not provided*

Contact Information

Preferred Contact


Name	Mark Magnuson
Institution	Vanderbilt University
Phone	615-322-7006
Email	mark.magnuson@vanderbilt.edu

Associated Publications

No publications associated

Comments

There are no comments for this entry.

 Login to add comments

[Home](#) · [Your Account](#) · [News & Events](#) · [Resources](#) · [Policies & Guidelines](#) · [About Us](#) · [FAQ](#) · [Site Map](#)

© 2002-2015 Beta Cell Biology Consortium - All Rights Reserved. [Terms of usage and disclaimer](#).

