

My Account

Login
Create Account

Resources

View All (813)
Adenoviruses (137)
Antibodies (175)
Bioimages (67)
Genomics Studies (145)
mESC Lines (68)
Mouse Strains (120)
Miscellaneous (46)
Protocols (55)
Research Data (4)
Resource Tags (389)
Visualization (9)

Research & Cores

Core Facilities (5)
Research Highlights (5)
Research Networks
Research Objectives

Information

About the BCBC
BCBC Events
Branding & Logos
Career Opportunities
Health
NIH hESC Registry
Policies & Guidelines
Member Publications
Research Programs
Research Investigators
Member Directory
Tutorials

Sst.rTTA.LCA - ES Cell Line RES4545**ESC Line Information**


Cell Line Name:	Sst.rTTA.LCA
Parental Cell Line:	TL-1
Background Strain:	129
Culturing Protocol:	Std_mESC_Culture.doc
Description:	This allele is designed to express a tet-inducible reverse-transactivator under control of the somatostatin promoter. Lox66 and Lox2272 sites are inserted flanking 500bp of Sst promoter region, Sst exons I and II and Sst poly A site. The rTTA-beta-globin poly A cassette replaces Sst coding exons (I and II). The allele can be used to drive tet-inducible expression in somatostatin expressing cells (Delta-cells). Additionally, the Lox66 and Lox2272 sites allow for manipulations of the flanked region by RMCE in the future.

Genetic Alterations

1) Targeted Mutagenesis	
Type of Allele	Cassette Acceptor
Targeted Gene	Somatostatin (Sst - NCBI GeneID:20604)
Targeted Allele	Sst.rTTA loxed cassette acceptor (Sst.rTTA ^{LCA})
Description of Targeting Vector	The targeting vector contains 7.3 kb 5' and a 3.6 kb 3' homology arms. Lox66 and Lox2272 sites are inserted flanking 500bp of Sst promoter region and exons I and II of the somatostatin gene. The rTTA gene, with beta-globin polyA site, is put in place of Sst exons I and II. The vector also contains FRT-flanked puTK-EM7 Neo selection double selection cassette. PuTK is used for positive selection for targeting events with puromycin and negative selection for RMCE events with ganciclovir. EM7-Neo is used for positive selection in bacteria during BAC recombineering process.
Targeting Vector Genbank File	pSst.rTTA.LCA.gb
Citations	Not Available


Associated Images

Image 1


**Description:**

This figure illustrates the gene targeting strategy of Sst, replacing exons I and II with the rTTA gene and beta-globin polyA. The vector also contains FRT-flanked puTK-EM7 Neo selection double selection cassette. PuTK is used for positive selection for targeting events with puromycin and negative selection for RMCE events with ganciclovir. EM7-Neo is used for positive selection in bacteria during BAC recombineering process.

Access Status

 This resource is publicly viewable.

Request this Resource

 Request from a repository


Primary contributor: [Magnuson Lab](#)

Co-contributed by:

- [BCBC Mouse / ES Cell Core](#)
- [Herrera Lab](#)

Resource Tags

embryonic, es, esc, Sst.rTTA.LCA, stem, TL-1


 Login to edit tags

 Read more about tags

Resource History & Actions

Approved on Oct 12, 2012

Last modified on Apr 20, 2015

 Login to edit or request an edit

Related resources**BCBC**

No matching resources

Other Consortia


No matching resources

Data courtesy of [dkCOIN](#). Only public resources are displayed.

Reference:
Not provided

Repositories

Magnuson Lab

 Request this resource

Stock #: Not provided
Availability Notes: Not provided

Contact Information

Preferred Contact


Name	Mark Magnuson
Institution	Vanderbilt University
Phone	615-322-7006
Email	mark.magnuson@vanderbilt.edu

Associated Publications

No publications associated

Comments

There are no comments for this entry.

 Login to add comments

[Home](#) · [Your Account](#) · [News & Events](#) · [Resources](#) · [Policies & Guidelines](#) · [About Us](#) · [FAQ](#) · [Site Map](#)

© 2002-2015 Beta Cell Biology Consortium - All Rights Reserved. [Terms of usage and disclaimer](#).

